

Childhood Cancer in the Vicinity of Nuclear Power Plants

The 2007 “KiKK”^{*} Study – IPPNW Physicians Issue Warning:

“Young children develop cancer more frequently when they live near nuclear power plants (NPP).

It has to be assumed that radioactive emissions from NPP stacks are indeed not as harmless as previously believed.

Now it is time to act.”

Ulm, January 11th 2008 Young children living near to German nuclear power plants develop cancer and leukaemia more frequently than children living further away from them. There is a 60% increased rate of cancer and approximately 120% of leukaemia. These are the findings of the “Epidemiological Study of Childhood Cancer in the Vicinity of Nuclear Power Plants” (KiKK Study), commissioned by the German Federal Office for Radiation Protection (BfS). Although the design of the study, carried out by the Mainz Cancer Registry, is generally held to be correct, interpretation of the study's findings is vigorously disputed by the authors¹. Indications of an increase in the incidence of childhood cancer near nuclear power plants have been found for over 20 years², but they have not as yet been taken sufficiently seriously. The correlation has been unequivocally confirmed by the KiKK study. **Now it is time to act.**

Background to the 2007 KiKK Study

The KiKK Study was called for in 2001 by IPPNW and the Ulm Physician's Initiative in a large-scale public relations campaign³, because several studies carried out by Dr. Alfred Körblein of the Munich Environmental Institute⁴, - including a study on NPPs in the Bavarian region initiated by IPPNW – had shown a significantly higher incidence of childhood cancer in the proximity of nuclear power plants. Only after massive pressure and over 10,000 letters of protest to the authorities and ministries did the Federal Office for Radiation Protection (BfS) accept the necessity for further studies⁵. The study was then commissioned by the BfS in 2003 to be carried out by the Mainz Cancer Registry⁶.

There had already been reports of significant increases in the levels of leukaemia around English nuclear installations in the 1980s. There were also sharp increases in rates of leukaemia around the reprocessing plants at Sellafield and La Hague.

An increased incidence of leukaemia found close to the Krümmel nuclear power plant caused much concern from the beginning of the 1990s onwards. Few studies on the subject were known, however, and most of those that existed showed nothing conspicuous in the vicinity of nuclear power plants – at least in the official versions:

1992⁷ and 1997⁸ – Two studies by the Mainz Cancer Registry (Director: Prof. Michaelis, Institute for Statistics and Documentation of the University of Mainz (IMSD)), covered the periods of 1980 to 1990 and 1980 to 1995 respectively. The childhood cancer rates in the vicinity of the 20 German nuclear installations (of which three were decommissioned nuclear power plants and two were research reactors) were examined. Main finding – nothing conspicuous⁹.

^{*} „KiKK“ stands for „Epidemiologische Studie zu Kinderkrebs in der Umgebung von Kernkraftwerken“ (Epidemiological Study of Childhood Cancer in the Vicinity of Nuclear Power Plants)

The “Michaelis” study has been consistently criticised since 1992 by Prof. Roland Scholz in numerous IPPNW^{10 11} and other publications^{12 13}. A renewed analysis of the data in the IMSD studies in 1998 by Dr. Körblein and Prof. Hoffmann¹⁴ showed that there was a significant increase in the rate of childhood cancer within a radius of 5 kilometres. The increase was to be found only when operational nuclear power plants were taken into account, not the decommissioned plants, nor the research reactors. The increase was only found amongst infants under the age of 5 years old.

Methodology and findings of the 2007 KiKK Study

The results of the KiKK study were published in December 2007 in the European Journal of Cancer¹⁵ and in the International Journal of Cancer¹⁶. The study covered all 16 large reactor locations where the 20 nuclear power plants in Germany were in operation during this period of time (period of study: 24 years, 1980 – 2003). Since the Lingen and Emsland locations are only two kilometres apart, they were combined into one study region. In the first part of the study a total of 1592 under-fives with cancer were compared to a control group of 4735 children. The distance between the children’s homes and the power plants was precisely determined to within 25 meters. The main questions posed by the study were: “Do children under five years old more frequently develop cancer when living near a nuclear power plant?” and “is there a negative distance trend?” (In other words: does the risk increase the nearer one lives to the plant?) The results showed not only a 60% increase in the cancer rate and a 117% increase in leukaemia in infants within the 5 kilometre radius, but also a significant increase in the risk of cancer and leukaemia the closer one lived to the nuclear power plant.

In the second part of the study, which covered a shorter period of time and a selection of diagnoses (leukaemia, lymphomas and tumours of the central nervous system), it was tested whether other risk factors (confounders) could have had any appreciable effect on the main result of the study – the negative distance trend. This proved not to be the case for any of the studied risk factors. The proximity of residence to the nuclear power plant remains the only plausible influencing factor.

On the discussion on the “small number” of cases

After the findings of the study were published in December 2007, the authors frequently emphasised appeasingly, that the study basically “only” dealt with a small number of cases of cancer. 37 cases were observed where 17 would have been expected statistically. This means that in a period of 24 years there was less than one additional leukaemia case a year. The 20 additional cases were only to be found within the 5 kilometre radius and were all cases of leukaemia. The reciprocal distance rule implicit in the study, however, adds up to a total of 127 additional cases amongst infants for the whole region under study.

Moreover, it can be assumed that such effects do not confine themselves to small children. Older children and adults could also be affected. However, the rates of cancer development amongst these groups have not yet been the subject of a comparable systematic study anywhere in the world.

It seems to me that the argument is significant that the KiKK study, in its methods and the questions it posed, was not set up to determine the exact number of additional cases of cancer at all. One can always find larger or smaller numbers of ill children according to the random selection of the size of the study area and using different distance rules. The latest KiKK study has a methodological strength, however, in testing the distance trend (which was also the main question posed by the study). This overcame the disadvantage of classically dividing the area into circular sections. But the KiKK study is inappropriate for determining the absolute number of cases. The authors’ reference to the small number of cases is obviously meant to soften the highly charged controversy over the results of the study. In any case, the study proves that there is an increased risk that correlates to the proximity to nuclear installations. That the absolute number of additional cancer cases is not higher is in part due to the fact that the area around nuclear power plants is usually thinly populated.

Controversial interpretation of the KiKK study – was it only coincidence?

The authors of the study were at first surprised by the results they had arrived at. They quickly pointed out that the raised levels of childhood cancer and leukaemia in the vicinity of nuclear power plants could not be explained by radioactive emissions. They claimed that the doses of radioactivity calculated to be in the vicinity of nuclear power plants are below the average dose from natural background radioactivity. Since this is not compatible with current radiobiological thinking, they did not rule out the possibility of coincidence as an explanation.

The findings of the 2007 KiKK study invalidate those of the previous studies by the Mainz Cancer Registry (IMSD 1992 and 1997). This should not really be a surprise, since Körblein had already pointed these effects out many times, as had Körblein and Hoffmann in their reanalysis of the IMSD study in 1998. For this reason, Körblein was strongly attacked by the Mainz Cancer Registry and accused of “data dredging”¹⁷. However, the 2007 KiKK study completely confirms the IMSD reanalysis of 1998. The authors belonging to the Mainz Cancer Registry have also admitted in the meantime that their earlier studies had already shown an increased cancer and leukaemia risk for infants living in close proximity to German nuclear power plants.

So what is the cause? “Coincidence” already has a long and sad tradition as an ultimate and helpless example of interpretation in radiological causality research. Let me remind you of the attempts made to explain the raised levels of childhood leukaemia in the area near the Krümmel power plant and Geesthacht nuclear research centre. Previous inexplicable clusters were given as the explanation for another inexplicable cluster. Was it once again simply coincidence? Yet coincidence as an explanatory model was clearly held to be improbable by the external expert’s group, commissioned by the BfS to supervise the drafting, the execution and evaluation of the KiKK study¹⁸. In referring to coincidence, the Mainz authors are ignoring the current state of research.

Already in the summer of 2007, a comprehensive meta-analysis by Baker et al. on leukaemia in children living near nuclear power plants¹⁹ caused a sensation. They examined data contained in a total of 17 international studies carried out in Germany, Spain, France, Japan and North America during the period between 1984 and 1999. Epidemiologists at the University of South Carolina discovered an enhanced risk of between 14 and 21% of developing leukaemia for children under nine years of age, depending on distance. All of the people examined under the age of 25 had an increased morbidity probability of about 7-10% and the rates of mortality were raised by 2-18%.

Correlation between the rate of morbidity, emission measurements, calculation model for radiation exposure and the biological effects of radionuclides

In Germany, children that are living near nuclear power plants develop cancer and leukaemia more frequently than those living further away. This has long been only a supposition, but has now been clearly proven and is officially accepted²⁰.

If emissions have been correctly measured by monitoring the areas surrounding nuclear installations, as has been claimed by both the NPP operators and the regulatory authorities, then either the currently accepted calculation models for determining radiation exposure of local residents are incorrect, or the biological effects of incorporated radionuclides have been badly underestimated, at least for young children or embryos.

The results of the KiKK study compel us to critically review not only the measurement of emissions by the operators but also the rules for calculating dose measurement and the risk models on which they are based. Any of these three steps could help solve the contradiction between the allegedly low doses and the severe effects referred to by the authors.

A separate inquiry into the boiling water reactor design type is also necessary

Boiling water reactors (BWR) have only one main cycle in their design. Pressure water reactors (PWR) have two separate main cycles, which means that BWRs have one less barrier holding back radioactive material from the surrounding area. The weak point can be found in the machine room of the NPP where highly radioactive hot steam is transported out of the reactor itself to the turbines.

In order to eliminate one individual location as the sole cause of the morbidity rate, the data in the study was assessed 16 times, each time excluding one location. In every case the exclusion did not change anything related to the main result of the study – a negative distance trend. It was not tested, however, whether there was a difference in risk levels when comparing BWR or PWR design types. This question could be answered easily enough using the existing study data.

There is enough evidence to show that the BWRs in Germany (currently Brunsbüttel, Krümmel, Phillipsburg 1, Isar 1, Gundremmingen B + C; in the past Würgassen) have higher levels of emissions. According to the annual reports of the government, environmental radioactivity and radiation exposure²¹ of BWRs are appreciably higher than those of PWRs, though within the currently accepted limits.

It is now time for us to act. The indications over many years that there are increased levels of morbidity near to NPPs have now been scientifically proven by the KiKK study.

No one can rule out the possibility of an increased risk for older children and adults living near NPPs. A systematic investigation of the type of the KiKK study has still to be carried out for these groups.

The previous mode of measuring emissions and reporting them needs to be put to the test. We can no longer rely on the information given by the NPP operating company. There needs to be official monitoring without any gaps and measured values must be made public.

Previous assumptions about radiation risk, and the emission limits for radiation that are based on these, need to be critically re-examined and adapted to current international research findings.

In addition, the data in the KiKK study should be separately assessed according to whether the location is a BWR or PWR.

We should primarily think about the people affected – the precautionary principle is long overdue.

Further cases of cancer near to NPPs have to be prevented. The only kind of reactor that does not present a cancer risk is a decommissioned reactor.

Reinhold Thiel, Spokesperson for the Ulm Physician's Initiative

Translated by Xanthe Hall, IPPNW

Internet links (in German)

www.ippnw-ulm.de

www.ippnw.de

www.alfred-koerblein.de

www.bfs.de

www.umweltinstitut.org

www.kinderkrebsregister.de

Acknowledgements

For feedback, thanks go to the following people: Dr. Winfrid Eisenberg, Dr. Veit Göller, Prof. Dr. Wolfgang Hoffmann, Raimund Kamm, Dr. Alfred Körblein, Prof. Dr. Dr. Edmund Lengfelder, Henrik Paulitz, Dr. Sebastian Pflugbeil, Prof. Roland Scholz und Karin Wurzbacher.

Thank you to the Federal Office of Radiation Protection for the constructive cooperation since July 2001, and for enabling and financing the KiKK study.

Thank you to the authors of the Cancer Registry at the University of Mainz for conducting the 2007 KiKK study, using a correct design (that was put forward by the external group of experts, that supervised the study).

Sources

- ¹ Unanimous Statement by the expert group commissioned by the German Federal Office for Radiation Protection on the KiKK Study (in German, no English version available), point 9, 5.12.2007, <http://www.bfs.de/de/kerntechnik/papiere/Expertengremium.html>
- ² Thiel R Kinderkrebs um Atomkraftwerke – Übersicht über nationale und internationale Studien, in IPPNW background information brochure „Nebenwirkungen der Atomenergie“, IPPNW 2002, http://www.ippnw-ulm.de/text_ueb-kinderkebs.htm
- ³ Thiel R Kinderkrebs um Atomkraftwerke, Chronologie eines 10-jährigen Engagements der IPPNW und der Ulmer Ärzteinitiative, 2.12.2007 <http://www.ippnw-ulm.de/071202-Chronologie-Kinderkrebs.pdf>
- ⁴ <http://www.alfred-koerblein.de/>
- ⁵ Protocol of meeting between BfS, IPPNW and the Munich Environmental Institute, Kassel, 11.07.2001 (Original in German, with author)
- ⁶ Thiel R, Paulitz H, Kinderkrebs um Atomkraftwerke, IPPNW Working group on nuclear energy press release, 19.03.2003, http://www.ippnw-ulm.de/text_krebs-akw0303.htm
- ⁷ Keller B, Haaf G, Kaatsch P, Michaelis J, Untersuchungen zur Häufigkeit von Krebserkrankungen im Kindesalter in der Umgebung westdeutscher kerntechnischer Anlagen 1980-1990, IMSD, technischer Bericht, Mainz, Institut für Medizinische Statistik und Dokumentation der Universität Mainz, 1992 – and: Michaelis J, Haaf G Kaatsch P, Keller B, Krebserkrankungen im Kindesalter, DÄ, Issue 30, 24.7.92, pp. 1386-1390
- ⁸ Kaletsch U, Meinert R, Miesner A, Hoisl M, Kaatsch P, Michaelis J, Epidemiologische Studien zum Auftreten von Leukämieerkrankungen bei Kindern in Deutschland, Bonn, Der Bundesminister für Umwelt, Naturschutz und Reaktorsicherheit, (1997 - Dr. Angela Merkel)
- ⁹ <http://www.kinderkrebsregister.de>
- ¹⁰ Scholz R, Weil nicht sein kann, was nicht sein darf, IPPNW-FORUM, Issue 8, December 1992, pp. 23-31
- ¹¹ Scholz R, Kernkraft und Kinderkrebs, in IPPNW-Studienreihe, Band 4 1st edition 1993, Bedrohung des Lebens durch radioaktive Strahlung, pp. 57-63
- ¹² Scholz R, Zwölf Anmerkungen zur IMSD-Studie, Strahlentelex, No. 130-131, 6th year, 4.6.1992, pp. 1-9
- ¹³ Scholz R, Weil nicht sein kann, was nicht sein darf, Kommentar zur Michaelis-Studie aus Mainz, in: Käufliche Wissenschaft, Experten im Dienst von Industrie und Politik, Bultmann/Schmithals (Ed.), Knauer-Verlag 1994
- ¹⁴ Koerblein A, Hoffmann W. Childhood leukaemia in the vicinity of German nuclear power plants, Medicine and Global Survival Vol.61(1999), pp. 18-23
- ¹⁵ Spix C et al, Case-control study on childhood cancer in the vicinity of nuclear power plants in Germany 1980-2003, Eur J Cancer (2007), doi:10.1016/j.ejca.2007.10.024
- ¹⁶ Kaatsch P et al, Leukaemia in young children living in the vicinity of German nuclear power plants, Int J Cancer (2007)
- ¹⁷ Statement by the Mainz Cancer Registry on their homepage of 19.6.2001, <http://www.kinderkrebsregister.de/stellungnahme0601.html>
- ¹⁸ Unanimous Statement by the expert group commissioned by the German Federal Office for Radiation Protection on the KiKK Study (in German, no English version available), point 10, 5.12.2007, <http://www.bfs.de/de/kerntechnik/papiere/Expertengremium.html>
- ¹⁹ Baker P.J., Hoel d.G. (2007), Meta-analysis of standardized incidence and mortality rates of childhood leukaemia in proximity to nuclear facilities, European Journal of Cancer Care, No. 16, pp. 355-363
- ²⁰ Weiss W, Background information on the KiKK Study, Federal Office for Radiation Protection, <http://www.bfs.de/en/kerntechnik/papiere/kikk.html>
- ²¹ Annual reports of the Federal German Government, Umweltradioaktivität und Strahlenbelastung, published on the site <http://dip.bundestag.de>. The report from 2001, 14/9995 <http://dip.bundestag.de/btd/14/099/1409995.pdf>, for instance, shows four times the amount of radiation in the spent air coming from NPP Gundremmingen that contains two BWRs.